



## ***Proteus* sp. – an opportunistic bacterial pathogen – classification, swarming growth, clinical significance and virulence factors**

ANTONI RÓZALSKI<sup>1\*</sup>, AGNIESZKA TORZEWSKA<sup>1</sup>, MAGDALENA MORYL<sup>1</sup>, IWONA KWIL<sup>1</sup>, AGNIESZKA MASZEWSKA<sup>1</sup>, KINGA OSTROWSKA<sup>1</sup>, DOMINIKA DRZEWIECKA<sup>2</sup>, AGNIESZKA ZABŁOTNI<sup>2</sup>, AGATA PALUSIAK<sup>2</sup>, MAŁGORZATA SIWIŃSKA<sup>2</sup>, PAWEŁ STĄCZEK<sup>3</sup>

<sup>1</sup> Department of Immunobiology of Bacteria

<sup>2</sup> Department of General Microbiology

<sup>3</sup> Department of Genetics of Microorganisms

Institute of Microbiology, Biotechnology and Immunology, University of Łódź, Banacha 12/16, 90-237 Łódź, Poland

\* E-mail: rozala@biol.uni.lodz.pl

### **ABSTRACT**

The genus *Proteus* belongs to the *Enterobacteriaceae* family, where it is placed in the tribe *Proteeae*, together with the genera *Morganella* and *Providencia*. Currently, the genus *Proteus* consists of five species: *P. mirabilis*, *P. vulgaris*, *P. penneri*, *P. hauseri* and *P. myxofaciens*, as well as three unnamed *Proteus* genomospecies. The most defining characteristic of *Proteus* bacteria is a swarming phenomenon, a multicellular differentiation process of short rods to elongated swarmer cells. It allows population of bacteria to migrate on solid surface. *Proteus* bacteria inhabit the environment and are also present in the intestines of humans and animals. These microorganisms under favorable conditions cause a number of infections including urinary tract infections (UTIs), wound infections, meningitis in neonates or infants and rheumatoid arthritis. Therefore, *Proteus* is known as a bacterial opportunistic pathogen. It causes complicated UTIs with a higher frequency, compared to other uropathogens. *Proteus* infections are accompanied by a formation of urinary stones, containing struvite and carbonate apatite. The virulence of *Proteus* rods has been related to several factors including fimbriae, flagella, enzymes (urease - hydrolyzing urea to CO<sub>2</sub> and NH<sub>3</sub>, proteases degrading antibodies, tissue matrix proteins and proteins of the complement system), iron acquisition systems and toxins: hemolysins, *Proteus* toxin agglutinin (Pta), as well as an endotoxin - lipopolysaccharide (LPS). *Proteus* rods form biofilm, particularly on the surface of urinary catheters, which can lead to serious consequences for patients. In this review we present factors involved in the regulation of swarming phenomenon, discuss the role of particular pathogenic features of *Proteus* spp., and characterize biofilm formation by these bacteria.

**KEY WORDS:** *Proteus*, pathogenicity, mechanisms of virulence

### **1. Classification and environmental distribution of *Proteus* rods**

The genus *Proteus*, which was described for the first time by Hauser in 1885, belongs to the *Enterobacteriaceae* family. In this family it is placed in the tribe *Proteeae*, together with the genera *Morganella* and

*Providencia*. *Proteus* rods are distinguishable from most other genera by their ability to swarm across agar surfaces of solid media. The differentiation of *Proteus* rods according to the results of biochemical tests is shown in

the table 1. The genus *Proteus* currently consists of five species: *P. mirabilis*, *P. vulgaris*, *P. penneri*, *P. hauseri* and *P. myxofaciens*, as well as three unnamed *Proteus* genomospecies. *Proteus myxofaciens* is the only *Proteus* species without any significance in the pathogenicity of humans, it has been isolated from living and dead larvae of the gypsy moth *Porteria dispar* (O'HARA *et al.* 2000, JANDA & ABBOT 2006). Recently

described results of the analysis of partial sequences of the *rpoB* gene encoding the  $\beta$ -subunit of RNA polymerase, led to the conclusion that a genus-level genetic distance exists between *P. myxofaciens* and the species of the *Proteus-Providencia* group. Classification of *P. myxofaciens* in a new genus *Cosenzaea* gen. nov. has therefore been proposed (GIAMMANCO *et al.* 2011).

**TABLE 1.** Biochemical tests used in differentiation within the genus *Proteus* (RÓŻALSKI & STĄCZEK 2010, 2011). + 90–100 % positive, - 0 – 9.9 % positive; S – susceptible, R – resistant, V – variable.

Test	<i>P. mirabilis</i>	<i>P. vulgaris</i>	<i>P. penneri</i>	<i>P. hauseri</i>	<i>P. myxofaciens</i>
Salicin fermentation	-	+	-	-	-
Maltose fermentation	-	+	+	+	+
D-Xylose fermentation	+	+	+	+	-
Esculin hydrolysis	-	+	-	-	-
Ornithine decarboxylase	+	-	-	-	-
Indole production	-	+	-	+	-
Chloramphenicol susceptibility	S	V	R	S	S

*Proteus* microorganisms are widely distributed in the natural environment, including polluted water, soil, and manure. Due to their proteolytic activity, the ability to hydrolyze urea to ammonia and carbon dioxide, as well as the oxidative deamination of amino acids, these bacteria are involved in

the decomposing of the organic matter of the animal origin. They are also present in the intestines of humans and animals. *P. mirabilis* was most frequently isolated from dogs, cows and birds, whereas *P. vulgaris* was most frequently isolated from pigs and cold-blooded vertebrates (JANDA & ABBOT 2006).

## 2. Swarming phenomenon


Swarming is a multicellular differentiation phenomenon that allows a population of bacteria to migrate on a solid surface in a coordinate manner. It is important in movement of *Proteus* rods to new locations and most probably helps them in the colonization macroorganisms. It involves cell-to-cell signaling and multicellular interactions and is connected with the possibility of morphological differentiation of bacteria depending on growth media. *Proteus* are

dimorphic bacteria, which in liquid media are motile, peritrichously flagellated short rods (1.0 to 2.0  $\mu\text{m}$  in length with 6–10 flagella). These bacteria are called swimmer cells. However, when transferred onto solid media these short rods change into elongated (20–80  $\mu\text{m}$  in length), hyperflagellated, multinucleated, nonseptated swarmer cells. The latter migrate out from the inoculation site as long as the population of swarmer cells is reduced on solid surfaces. Then, the

consolidation process takes place. In this period of swarming growth, the long rods disintegrate to short bacteria. The processes of differentiation and dedifferentiation of *Proteus* bacteria are cyclic. It results in the formation of characteristic rings of bacterial growth on the agar plate (fig. 1) (VERSTRAETEN *et al.* 2008, MORGENSTEIN *et al.* 2010, RÓŻAŁSKI & STĄCZEK 2010).

Differentiation of swimmer cells into swarmer cells is induced by the contact of bacteria with a solid surface and the inhibition of flagellar rotation. It was found that the addition of thickening agents to liquid media resulted in differentiation of short rods to swarmer cells. A similar effect was obtained when anti-flagellar antibodies were added to liquid medium (BELAS & SUVANASUTHI 2005). Mutations in gene *flaA* encoding flagellin, as well in genes involved in flagellum assembly led to the inhibition of *Proteus* short rods differentiation (BELAS 1994). Lipopolysaccharide (LPS), a surface component of these bacteria, seems to be important in the swarming phenomenon, however, its exact role has not been

described. Mutation in the *wzz* region encoding O-antigen chain, as well as in genes *waaD* and *waaC* required for the core region of LPS are impaired in swarming growth (MORGENSTEIN *et al.* 2010). Swarmer cells translocation is facilitated by an extracellular acidic polysaccharide designated as Cmf (colony migration factor), which acts as a lubricant, reducing surface friction (GYGI *et al.* 1995). This capsular polysaccharide is structurally identical with the O-specific part of *Proteus* LPS serogroup O6 (KNIREL *et al.* 2011). Most likely, *Proteus* O-specific polysaccharides are important for the swarming. It was found that the Re mutant of *P. mirabilis* R45 producing LPS having only the lipid A and Kdo region was unable to swarm. The Ra mutant of *P. mirabilis* R110 containing lipid A and a complete core region expressed only a limited ability for migration on the solid medium, whereas most of S-forms of *P. mirabilis*, *P. vulgaris* and *P. penneri* could swarm vigorously (BABICKA 2001, KWIL 2003).



**Figure 1.** Swarming growth of three strains *P. mirabilis* strains 34/57 (A), 14/57 (B) and 26/57 (C) on a agar plate (KWIL data not published).

Swarming growth is characteristic for population of bacteria, thus it is dependent on multicellular interactions and cell to cell signaling in the phenomenon described as quorum sensing. It was found that

extracellular signal AI-2 (autoinducer-2) does not play a role in swarming. Cyclic dipeptides were also shown as *P. mirabilis* signaling molecules out of importance in swarming (RATHER 2005). Other signal molecules

*N*-acyl homoserine lactones crucial in quorum sensing are not produced by *P. mirabilis* (MORGENSTEIN *et al.* 2010). Nevertheless, it was shown that *N*-acyl homoserine lactones added to the media are able to accelerate migration of some *P. mirabilis* strains (STAŃKOWSKA *et al.* 2008). In earlier studies, glutamine was found to be able to initiate swarming on minimal media (ALLISON *et al.* 1993). Swarming growth is also influenced by fatty acids. Oleic acid stimulates, whereas lauric and myristic acids inhibit this phenomenon, respectively (LIAW *et al.* 2004). The mechanism by which glutamine and fatty acids play a role in the swarming process as signals for cell to cell communication is currently unknown. In swarmer cell differentiation putrescine was described as a biochemical signal, however, the mechanism by which this amine regulates swarming is not clear. Most probably putrescine can form a complex with galacturonic acid which is component of CPS. This suggests a possibility that putrescine complexed with cell surface polysaccharide acts as a signal factor or putrescine regulates expression of CPS (RATHER 2005, MORGENSTEIN *et al.* 2010).

A number of factors playing a role in swarming regulations have been recognized (DANIELS *et al.* 2004, RATHER 2005, VERSTRAETEN *et al.* 2008, MORGENSTEIN *et al.* 2010). The most important gene involved in upregulation of flagellin production is *flhDC*, class 1 gene of flagellar regulon. This gene encodes FlhD2C2 complex, a heterotetrameric transcriptional regulator, which also regulates the expression of additional genes required for swarmer cells differentiation. The expression of *flhDC* increases 10-fold during initiation of swarmer cells differentiation and it is influenced by a variety of environmental factors and regulating genes. FlhD2C2 activates class 2 genes coding basal body and hook proteins of flagella, as well as  $\sigma_{28}$  factor, which activates class 3 genes required for synthesis of flagellin and flagellum assembly. FlhD2C2 activity is regulated by two identified factors DisA (decarboxylase inhibitor of swarming) and Lon protease (ATP-dependent protease).

DisA most probably inhibits the assembly of heterotetramer or its binding to DNA (MORGENSTEIN *et al.* 2010).

*Proteus* has 16 predicted two-component systems – TCS (PEARSON *et al.* 2008). TCS consists of sensor kinase, which activates response regulator DNA – a binding protein controlling gene expression. Two of TCS – Rcs and Rpp are involved in swarming regulation (MORGENSTEIN *et al.* 2010). Rcs system contains RcsC sensor kinase, RcsB response regulator, RcsD intermediate transferring phosphate to the RcsB and RcsF which is an outer membrane lipoprotein. Mutation in the genes coding Rcs system in *P. mirabilis* lead to the hyperswarming phenotype, most probably due to an increased expression of the *flhDC*. *P. mirabilis* Rcs mutant grows in liquid media as elongated cells. The Rpp system consists of RppA response regulator and RppB histidine sensor kinase like protein. Mutation in the Rpp system in *P. mirabilis* also results in a hyperswarming phenotype.

*flhDC* expression is also regulated by several proteins including Umo, MrpJ, WosA, Lrp and RsmA. Umo A-D proteins (umo – upregulated expression of the *flhDC* master operon) are located in the cell membrane and periplasm. Mutant defective in production of UmoD protein is not able to swarm, however, such effect was not noticed in the case of UmoA and UmoC (DUFOUR *et al.* 1998).

*mrpJ* gene is located in the *mrp* operon encoding MR/P fimbriae, which are required for sessile lifestyle of bacteria (NIELUBOWICZ & MOBLEY 2010). It is opposite to the swimming or swarming phenotype of bacteria. The existence of bacteria in one of two forms is regulated by *mrpJ* gene product – MrpJ protein, a transcriptional regulator. It binds the *flhDC* promoter, which causes its repression and results in loss of motility of bacteria (PEARSON & MOBLEY 2008).

*wosA* gene coding WosA protein, is overexpressed in strains exhibiting hyperswarming phenotype (*wos* – wild type onset superswarming). WosA overexpression increases the expression of *flhDC* and leads to a differentiation of swarmer cells in liquid

media. The hyperswarming *wosA* bacteria move quickly and spend less time in the consolidation phase. *wosA* expression is growth phase dependent, as well as it is partially dependent upon the expression of *flaA* gene encoding the flagellar filament (HATT & RATHER 2008).

Leucine responsive protein (Lrp) is a transcriptional global regulator involved in regulating different processes including amino acids synthesis, peptide transport, and pilin biogenesis (RATHER 2005, MORGENSTEIN *et al.* 2010). In *P. mirabilis* strains Lrp plays a role in the regulation of the swarming phenomenon, most probably its action leads to the inhibition of hyperflagellation. Mutation in *lrp* results in a decrease in the *flhDC*, *flaA* and *hpm* (hemolysin) expression and, in consequence, a nonswarming phenotype (HAY *et al.* 1997). Lrp is well known to respond to leucine and to a lesser extent to alanine. Recently, it was found that it also responds to methionine and is sensitive to isoleucine, histidine and tyrosine (HART & BLUMENTHAL 2010). In additional studies HART *et al.* (2011) showed that recognition of DNA by the helix-turn-helix motif of Lrp is modulated by the N-terminal tail.

RsmA – repressor of secondary metabolites is a member of regulatory systems involved in the expression control of genes which code factors committed to stationary growth processes. This protein stabilizes mRNA. In *P. mirabilis* RsmA is 6.8 kDa protein containing 62 amino acids. Increasing its expression results in the inhibition of swarmer cells differentiation and production

of hemolysin, protease, urease and flagellin (LIAW *et al.* 2003). Overexpressing of *rsmA* led to repression of *flhDC* (MORGENSTEIN *et al.* 2010).

The role of the swarming phenomenon in the pathogenicity of *Proteus* bacteria (see below) is till now unclear. It was shown that swarmer cells demonstrate higher production of urease, HpmA hemolysin and IgA metalloprotease ZapA, as well as flagellin synthesis, as compared to swimmer cells (RATHER 2005, JACOBSEN *et al.* 2008). There is some controversy concerning invasiveness of swarmer cells. ALLISON *et al.* 1992 showed *in vitro* that this morphotype could invade uroepithelial cells, whereas nonflagellated and nonswarming forms were noninvasive. Other authors (RÓŻALSKI *et al.* 1986, CHIPPENDALE *et al.* 1994) showed the short rods as invasive forms, rather than swarmer cells. These results were confirmed by JANSEN *et al.* (2003) in a mouse model of ascending UTI. These authors have found that the predominant cell type in the urinary tract are short swimmer cells but not elongated swarmer cells. Recently, FUJIHARA *et al.* (2011) suggested that *P. mirabilis* cells differentiate into hyperflagellated and multinucleated swarmer cells in acidic pH of the host's urine and differentiate back into swimmer forms, when the urinary pH is increased and it is alkaline after urease action (see below). In acidic condition swarmer cells exhibit higher cytotoxicity against T24 line. In alkaline condition *P. mirabilis* showed few elongated cells with high number of flagella and cytotoxic activity.

### 3. Pathogenicity

*Proteus* rods are opportunistic bacterial pathogens which under favorable conditions cause urinary tract infections (UTIs), wound infections, meningitis in neonates or infants and rheumatoid arthritis (O'HARA *et al.* 2000, JANDA & ABBOT 2006). KALRA *et al.* (2011) reviewed endocarditis due to *Proteus* species and OKIMOTO *et al.* (2010) reported *P. mirabilis* pneumonia. Brain abscesses during *P. vulgaris* bacteremia were described

by BLOCH *et al.* (2010). However, it should be stressed that *Proteus* bacteria cause UTIs with higher frequency. This type of infections is classified as uncomplicated or complicated. Uncomplicated infections occur in patients who are otherwise considered healthy, whereas complicated infections usually take place in patients with an urinary catheter in place or with structural and/or functional abnormalities in the urinary tract, suffering

from another illness, immunocompromised, as well as after surgical intervention in the urogenital system. It was found that *Escherichia coli* is a common cause of uncomplicated infections. Complicated UTIs might be polymicrobial and are usually caused by Gram-negative bacteria *Proteus* spp., *Providencia stuartii*, *Morganella morganii*, *E. coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* as well as some Gram-positive bacteria (NIELUBOWICZ & MOBLEY 2010). *Proteus* rods can cause hematogenous infections and ascending infections, however, the latter are more common for these microorganisms. The ascending infections comprise colonization subsequently urethra, bladder, ureters and kidneys. *P. mirabilis* causes UTIs with the highest frequency among all *Proteus* species. These bacteria are a common cause of complicated UTIs with the frequency of 12 % and catheter associated bacteriuria (WARREN 1996, JACOBSEN *et al.* 2008). They also cause uncomplicated lower urinary tract infections, however less frequently. CHLABICZ *et al.* (2011) have found in Poland uncomplicated *P. mirabilis* infections with the frequency of 3.4 %.

The virulence of *Proteus* rods has been related to several factors including fimbriae; flagella; enzymes: urease, hydrolyzing urea to CO<sub>2</sub> and NH<sub>3</sub>; proteases degrading antibodies, tissue matrix proteins and proteins of complement system; iron acquisition systems and toxins: hemolysins, *Proteus* toxin agglutinin (Pta), as well as endotoxin - lipopolysaccharide (LPS) (table 2) (COKER *et al.* 2000, MOBLEY 1996, RÓZALSKI *et al.* 1997, RÓZALSKI 2002, NIELUBOWICZ & MOBLEY 2010). BEVERIDGE (1999) postulated the possible role of outer membrane vesicles (OMVs) released from the *P. mirabilis* surface, as an alternative route of delivery of some enzymes, as well as some virulence factors.

PEARSON *et al.* (2011) using microarrays analyzed *P. mirabilis* gene expression *in vivo* from experimentally infected mice and showed 471 gene upregulated and 82 downregulated in comparison to the *in vitro* broth culture. Upregulated genes encoded

among others mannose-resistant *Proteus*-like fimbriae, urease, and iron uptake systems. Genes encoding flagella were downregulated. Earlier studies (FLANNERY *et al.* 2000) showed the identification of 94-kb PAI (pathogenicity island) designated ICEPm1 (integrative and conjugative elements *P. mirabilis* 1). This PAI is common to *P. mirabilis*, *P. stuartii* and *M. morganii* and carries virulence factors including *Proteus* toxic agglutinin (see below) and the high pathogenicity island of *Yersinia* spp.

The *P. mirabilis* genome predicts 17 distinct fimbrial operons and 13 additional orphan fimbrial genes not associated with complete operons (PEARSON *et al.* 2008, NIELUBOWICZ & MOBLEY 2010). The biogenesis and/or the biological role were studied for six types of fimbriae and agglutinins: MR/P (mannose resistant *Proteus* like fimbriae), MR/K (mannose resistant *Klebsiella* hemagglutinins), PMF (*Proteus mirabilis* fimbriae), NAF (nonagglutinating fimbriae), ATF (ambient temperature fimbriae) and *P. mirabilis* P-like fimbriae (fimbriae, which are similar to *E. coli* P pili) (COKER *et al.* 2000, ROCHA *et al.* 2007b, JACOBSEN *et al.* 2008).

The most important *P. mirabilis* fimbriae seem to be MR/P. This type of fimbriae is encoded by *mrp* operon containing 10 genes located on bacterial chromosome. The main structural subunit of these fimbriae is the MrpA protein. The MrpC protein is an outer membrane platform, MrpD plays a role of a chaperon protein and MrpH is an adhesin located at the fimbrial tip (BAHRANI *et al.* 1991, LI *et al.* 1999, COKER *et al.* 2000). MR/P/ fimbriae undergo phase variation. It was found that the MrpI protein acts as *P. mirabilis* recombinase controlling “on/off” orientation of the *mrp* promoter. The fimbrial genes are transcribed and MR/P fimbriae are synthesized when the promoter is in “on” orientation (LI *et al.* 2002a, NIELUBOWICZ & MOBLEY 2010). In the “off” orientation the production of fimbriae is stopped. Expression of MR/P fimbriae is increased under oxygen limitation (LANE *et al.* 2009). These fimbriae contribute to biofilm formation and facilitate

colonization of upper urinary tract and were found more often on bacterial strains which cause pyelonephritis (JANSEN *et al.* 2004). MR/P fimbriae are involved in aggregative adherence to HEp-2 cells (ROCHA *et al.* 2007a). MR/P fimbriae elicit a strong immune response during infection. It is directed against the MrpA protein. It was found that immunization of mice with the attenuated strain of *Salmonella* Typhimurium expressing MrpA-TetC fusion protein led to an increase in serum level of IgG antibodies against MrpA and significantly reduced *P. mirabilis* colonization of bladder and kidney. TetC is a non-toxic highly immunogenic fragment of the tetanus toxin (SCAVONE *et al.* 2011). The presence of *mrp* operon was shown not only in *P. mirabilis* strains but also in other species of the *Proteus* genus (BABICKA 2001, KWIL, 2002, 2003, KWIL *et al.* data not published).

Other than MR/P fimbriae are known to a lesser extent. MR/K hemagglutinins were found more frequently on *P. penneri* strains than on *P. mirabilis* strains and most probably they facilitated the adherence of bacteria to the urinary catheters (JACOBSEN *et al.* 2008). The exact virulence role of PMF fimbriae is unclear (NIELUBOWICZ & MOBLEY 2010). ZUNINO *et al.* (2007) by using *P. mirabilis pmfA/mrpA-D* mutant showed the specific and additive roles of MR/P and PMF fimbriae in UTI caused by these bacteria. NAF fimbriae, earlier known as UCA fimbriae, allow bacteria to colonize uroepithelial cells (WRAY *et al.* 1986). It was found that NAF/UCA binds the following receptors: asialo-GM<sub>1</sub>, asialo-GM<sub>3</sub>, lactosyl ceramide and galectin (LEE *et al.* 2000, ALTMAN *et al.* 2001). ATF fimbriae are not important in the pathogenicity of human beings (ZUNINO *et al.* 2000). PMP fimbriae were found on canine UTI isolates (BIJLSMA *et al.* 1995), however genes coding this type of fimbriae were found in human clinical isolates (PEARSON *et al.* 2008).

There are conflicting results concerning the importance of flagella in the infection. Since flagella mediate motility of bacteria, they seem to be required for virulence of uropathogens, particularly during ascending

infection. Indeed, it was shown *in vivo* that flagella-negative mutant of *P. mirabilis* is less virulent, compared to the wild strain, which suggests an important role of these bacterial surface structures in the pathogenicity (MOBLEY *et al.* 1996). However, it must be stressed, that lack of flagella on *P. mirabilis* human isolate was found (ZUNINO *et al.* 1994). Flagella play a role of H antigen and are strongly immunogenic. Bacteria can avoid immune response of human organism due to the antigenic variation process. This phenomenon is based on flagellin genes rearrangement. Flagellar antigenic variations allow bacteria to evade the action of secretory IgA antibodies directed against these organelles (BELAS 1994).

*P. mirabilis* evades the immune system response during infection thanks to the production of sIgA protease. The best characterized is 54-kDa secreted metalloprotease ZapA, mirabilysin belonging to the serralyisin family (LOOMES *et al.* 1990). ZapA possesses a broad spectrum of proteolytic specificity, including the activity against antibodies of sIgA, IgA2 and IgG, proteins of the complement system, cell matrix proteins, cytoskeletal proteins, as well as antimicrobial defense components of human innate system peptides hBD1 and cathelicidin LL-37 (BELAS *et al.* 2004, JACOBSEN *et al.* 2008). ZapA was found to be an important virulence factor in a rat model of *P. mirabilis* induced acute and chronic prostatitis (PHAN *et al.* 2008). Recently, CARSON *et al.* (2011) described *N*-alpha mercaptoamide dipeptides, which can be used as inhibitors of ZapA. KWIL *et al.* (2011) have shown the presence of the *zapA* operon in *P. vulgaris* and *P. penneri* strains.

It was shown that amino acid deaminase (Aad) produces  $\alpha$ -keto acids which serves as siderophores involved in iron acquisition (DRECHSEL *et al.* 1993, MASSAD *et al.* 1995). Recently, HIMPSI *et al.* (2010) described proteobactin and yersiniabactin related siderophore function in *P. mirabilis*. Zinc, another biologically important element needed for bacterial growth, is utilized by *P. mirabilis* by use of ZnuABC high affinity

transport system (NIELUBOWICZ & MOBLEY 2010). *P. mirabilis* also posses high-affinity phosphate transporter system Pst. It is postulated that this system is important in *P. mirabilis* virulence (JACOBSEN *et al.* 2008).

Urease is an urea amidohydrolase which catalyses the hydrolysis of urea to ammonia and carbamate, then digested to carbone dioxide and second molecule of ammonia (MOBLEY *et al.* 1995). This process results in elevation of pH and non-physiological alkalization of urine, which in turn induces

the precipitation of magnesium and calcium ions, normally soluble in slightly acidic urine, and the formation of urinary stones, containing struvite and carbonate apatite (RÓŻALSKI *et al.* 1997, JACOBSEN *et al.* 2008, NIELUBOWICZ & MOBLEY 2010). The carbonate apatite forms an amorphous precipitate opposite to struvite, which forms crystals of defined morphology. It was found that struvite exhibits polar properties (PRYWER & TORZEWSKA 2010, ROMANOWSKI *et al.* 2010).

**Table 2.** Virulence factors (VP), biological features (BF) and types of growth (TG) of *Proteus* rods in relation to the pathogenicity.

VP/BF/TG	Specific data	Contribution to the pathogenicity
Fimbriae	<i>P. mirabilis</i> genome predicts 17 distinct fimbrial operons. Six types of fimbriae/agglutinins were identified	Adherence of bacteria to the epithelial tissue Contribution to the biofilm formation
Flagella	H-antigen	Ascension of bacteria in urinary tract. Elicitation of immunoresponce which can be avoid due to the antigenic variation
Invassiveness	Studied <i>in vitro</i>	Penetration and internalization of bacteria into host cells
Proteases	IgA and IgG proteases, which are also able to degrade other types of biologically important proteins	Diminish or elimination of phagocytosis
Amino acid deaminase	Production of alpha-keto acids which serve as siderophores	Iron acquisition
Proteobactin and yersiniabactin - related siderophore		
ZnuABC	High affinity zinc transport system	Zinc utilization
Pst	High affinity phosphate transporter	Phosphate accumulation
Urease	Enzyme hydrolyzing urea to NH <sub>3</sub> and CO <sub>2</sub>	Elevation of urine pH level, resulting in stones formation. Cytotoxicity against urinary epithelial cells
Protein toxins:		
HmpA	Cell-bound hemolysin	Pore forming toxins mediating cytotoxicity
HlyA	Cell free hemolysin, RTX protein	
Pta	Cytotoxin and agglutinin	Cytotoxicity and autoagglutination of bacteria
CPS	Most probably structurally identical with O-specific part of LPS	Formation of biofilm. Importance in formation of urinary stones. Facilitation of swarmer cells translocation
LPS	Endotoxin	Endotoxicity. Resistance against bactericidal action of the serum. Production of glycocalyx important in biofilm formation
OMVs	Outer membrane vesicles containing periplasmic constituents	Transport of virulence factors to the host tissues
Swarming growth	Differentiation of short rods to swarmer cells	Not fully understood
Biofilm formation	Crystalline biofilm, particularly formed on the surface of urinary catheters	Protection of bacteria against action of antibiotics and immune mechanisms



The structural and elastic properties of struvite crystals were studied by PIECHOTA *et al.* (2011). Formation and morphology of struvite crystals depend on the pH of urine. PRYWER and TORZEWSKA (2009) have found that for pH in the range 7.5–9.0 struvite crystals take single and hemimorphic morphology. At higher pH many twin and dendritic crystals have been shown. DUMANSKI *et al.* (1994) has revealed that *Proteus* capsular polysaccharide structurally identical to O-specific polysaccharide of LPS accelerates crystallization process. TORZEWSKA *et al.* (2003) confirmed this observations and showed that urinary stone formation due to the urease activity, may be modified by surface O-antigens being a part of lipopolysaccharides (LPS). The sugar composition of these polysaccharides may either enhance or inhibit the crystallization of struvite and carbonate apatite depending on their chemical structure and ability to bind cations. Bacteria present within urinary stones are protected from host defense mechanisms and antibiotic treatment (LI *et al.* 2002b). Therefore, several phytotherapeutic compounds have been investigated to prevent urinary stone formation. For example curcumin an yellow-orange pigment non toxic to humans or animals extracted from the roots of turmeric (*Curcuma longa*) was shown to inhibit urease activity and consequently decrease the struvite growth. It was found *in vitro* experiment of struvite formation in artificial urine that the addition of curcumin increases the induction time and decreases the efficiency of formation of these crystals (TORZEWSKA *et al.* 2010, PRYWER & TORZEWSKA 2010, 2012). Moreover, curcumin also inhibits biofilm formation and block swarming growth (TORZEWSKA *et al.* 2010). Recently, MALIC *et al.* (2011) reported on the development of an early warning sensor for encrustation of urinary catheters in case of *Proteus* infection.

It was shown that the high level of ammonia, a product of urease action also results in damage of the urinary tract epithelium. Toxic ammonia damages bladder surface glycosaminoglycans important in the

prevention of bacterial adherence to the uroepithelium (JACOBSEN *et al.* 2008). *In vitro* studies performed by TORZEWSKA and RÓZALSKI (2009) showed that glycosaminoglycans have no inhibitory effect on infection-induced crystallization.

In a mouse model of ascending UTI it was found that the urease negative mutant of *P. mirabilis* was less virulent as compared to the wild strains, which suggests the important role of this enzyme in infection caused by this bacterium (JONES *et al.* 1990). Therefore, investigation of different substances which can be used as urease inhibitors is being performed. Recently, BERLICKI *et al.* (2011) described N-substituted amino-methane-phosphonic and aminomethane-P-methylphosphonic acids as effective inhibitors of *P. mirabilis* urease.


*Proteus* rods produce three types of cytolytins, HpmA and HlyA hemolysins and Pta cytotoxic agglutinin. HpmA is the 166 kDa protein, calcium-independent cell-bound hemolysin secreted by the bacteria and activated in the process mediated by HpmB (UPHOFF & WELCH 1990). This hemolysin belongs to the pore forming toxins family and it is cytotoxic against a number of cells (SWIHART & WELCH 1990). HlyA hemolysin is produced by *P. vulgaris*, *P. penneri* and *P. hauseri* strains (KORONAKIS *et al.* 1987, RÓZALSKI & KOTELKO 1987, RÓZALSKI & STĄCZEK 2011). It is a strongly cytotoxic, calcium dependent 110 kDa extracellular toxin belonging to the RTX family of proteins (RTX - repeat in toxin). Pta (*Proteus* toxic agglutinin), an autotransporter with subtilase-like serine protease activity exhibits bifunctional activity. It is both cytotoxin and agglutinin, which mediates autoagglutination of bacteria and cytotoxicity against human bladder and kidney cells. The vaccination of mice with Pta results in protection against *P. mirabilis* UTI (ALAMURI & MOBLEY 2008).

Lipopolysaccharide is composed of three genetically and structurally distinct regions: O-specific chain (O-antigen, O-specific polysaccharide), the core oligosaccharide and lipid A, which anchors the LPS molecule to the bacterial outer membrane (RAETZ &


WHITFIELD 2002). LPS containing all these three regions is produced by smooth forms of bacteria. Rough strains synthesize LPS containing lipid A and the whole core region or only its part. All three regions of *Proteus* LPS have been studied (RÓŻALSKI 2002, 2004, 2008, RÓŻALSKI *et al.* 2002). The differences in the structure of O-antigens serve as a basis for the serological classification of *Proteus* strains. The serological classification scheme currently consists of 78 serogroups (DRZEWIECKA *et al.* 2004, 2008, 2010, DRZEWIECKA & SIDORCZYK 2005, KNIREL *et al.* 2011). The prevalence of particular *Proteus* O-serogroups among clinical isolates has been changing over the time. Previously, clinical isolates

were found to belong to serogroups O3, O10, O11, O13, O23, O27 and O30. Recently, KACA *et al.* (2011) studying 123 clinical isolates from Sweden and Poland showed that none of serogroups was predominant. DRZEWIECKA *et al.* 2010 described seven *P. mirabilis* strains representing serogroup O78 which can be classified in one bacterial clone widespread in hospitals, most probably due to the nosocomial infection and autoinfections. The results of detailed studies of *Proteus* O-specific polysaccharides are presented in the review by KNIREL *et al.* 2011. WANG *et al.* (2010) identified five putative *Proteus* O-antigen gene clusters and *Proteus* specific genes of strains classified to five serogroups.

**A**



**B**



**Figure 2.** Structure of the O-specific repeating unit (A) and core oligosaccharide obtained by mild acid degradation (B) of lipopolysaccharide of *Proteus mirabilis* genomic strain HI 4320. Alt - altruronic acid (KNIREL *et al.* 2011, VINOGRADOV 2011).

The core region of *Proteus* lipopolysaccharides was studied by use of rough mutants or smooth forms classified into different serogroups (RADZIEJEWSKA-LEBRECHT *et al.* 1989, VINOGRADOV *et al.* 1994, VINOGRADOV *et al.* 2002). The structural diversity of the core is characteristic for *Proteus* spp. and makes it different from

*E. coli* and *Salmonella* (HOLST 1999, VINOGRADOV *et al.* 2002). The *Proteus* core region is composed of two parts – an inner part, common for several number of strains and second, an outer part, which is characterized by structural variability from strain to strain. The common part is not identical in all *Proteus* strains and is

subdivided into three forms known as glycoforms I-III. The outer part of the core region (outer core) contains an oligosaccharide characteristic for particular *Proteus* strains (VINOGRADOV *et al.* 2002, RÓZALSKI 2008). The structure of the core part of *P. mirabilis* genomic strain HI4320 (formerly classified into serogroup O10) was shown by VINOGRADOV (2011) (fig. 2). The epitope specificity of the core region of *P. penneri* strains were studied by PALUSIAK and SIDORCZYK (2010). They have identified seven epitopes responsible for cross reactivity of LPS of these bacteria with appropriate antisera. The identification of genes required for the biosynthesis of the core region of *P. mirabilis* were described by AQUILINI *et al.* (2010).

LPS, released from pathogenic bacteria during infection induces a spectrum of biological activities. Therefore, LPS is also known as an endotoxin. The mechanism of biological action of the endotoxin is common to most bacteria. Its biological domain is lipid A (ŁUKASIEWICZ & ŁUGOWSKI 2003). *Proteus* lipid A contains glucosamine disaccharide substituted with phosphate residues and fatty acids. It also contains 4-amino-4-deoxy-L-arabinopyranose (L-Arap4N), which quantitatively substitutes the ester-linked phosphate residue of the glucosamine backbone (SIDORCZYK *et al.* 1983).

*Proteus* LPS is associated with a broad spectrum of biological activities, and with interactions with bacterial or eukaryotic cells (RÓZALSKI 2008). LPS of S-forms of bacteria contributes to their resistance against bactericidal action of the serum (KACA & UJAZDA 1998, MIELNIK *et al.* 2004, BABICKA 2001, KWIL 2003, KACA *et al.* 2009). The O-specific polysaccharide chain exposed outside bacteria is involved in glycocalyx formation. Glycocalyx enables bacteria to grow in a form of biofilm on a solid surface.

Bacterial biofilm is defined as a matrix-enclosed bacterial population adhering to the surfaces. Bacteria enclosed in a glycocalyx capsule are protected against the action of antibodies, as well as against other immune

mechanisms (DONLAN & COSTERTON 2002, STICKLER & FENLEY 2010). *Proteus* bacteria form a crystalline biofilm particularly on the surface of catheters (STICKLER 2008). It contains calcium phosphate and magnesium phosphate crystals formed due to the action of urease. JONES *et al.* (2007) reported that the structure of *P. mirabilis* biofilm depends on the environment (artificial urine, laboratory media) in which it occurs. Biofilms in Luria-Bertani (LB) broth formed a mushroom-like architecture, while biofilms grown in urine formed a flat layer. MORYL *et al.* (2008) studied the effect of environmental factors such as glucose, acidic pH, albumin and some antimicrobial agents: ciprofloxacin and methylene blue on the formation of *P. mirabilis* biofilm. The most inhibiting effect was observed after incubation of bacteria with albumin and in acidic growth media. KWIECINSKA-PIRÓG (2010) studied the formation of biofilm on different surfaces, and has found silicone latex surface to be better than polychloride vinyl and polypropylene surfaces for *P. mirabilis* growth in biofilm. Octenisept R an antiseptic products occurred to be active against these bacteria growing in biofilm. Prevention and eradication of *P. mirabilis* biofilm by use of lytic bacteriophages was shown by CARSON *et al.* (2010). It was also found that two quorum sensing antagonists p-nitrophenyl glycerol (PNPG) and tannic acid inhibit the formation of *P. mirabilis* biofilm in artificial urine (JONES *et al.* 2009). *P. mirabilis* biofilm formation on glass and polystyrene can also be inhibited by extract of *Ibucella lutea*, a native plant of America (SOSA & ZUNINO 2009). MORYL (2010) observed inhibition of biofilm formation by *P. mirabilis* growing together with *Escherichia coli* or *Providencia stuartii*. STAŃKOWSKA *et al.* (2011) showed a significant influence of *N*-butanoyl homoserine lactone (BHL) on biofilm formation by *P. mirabilis* strain belonging to serogroup O18. Recently, SCHLAPP *et al.* (2011) excluded the contribution of swarmer cells in biofilm formation during the 7-day incubation of bacteria in LB broth on glass coverslips. These authors showed that

*P. mirabilis* biofilm formation is a five-stage process: i) reversible adhesion to the solid surface with slow growth of bacteria, presence of elongated cells and absence of extracellular matrix, ii) irreversible bacterial adhesion, beginning of production of

extracellular materials and decreasing of elongation, iii) acceleration of bacterial growth, further decrease in elongation and halting production of extracellular material, iv) maturation of biofilm, v) detachment of bacterial cells and dispersion of biofilm.

#### 4. Summary

*Proteus* bacteria inhabit the environment and they are also present in the intestines of humans and animals. These microorganisms under favorable conditions cause a number of infections including urinary tract infections (UTIs), wound infections, meningitis in neonates or infants and rheumatoid arthritis. *Proteus* opportunistic pathogens express virulence factors associated with adherence, motility, immunoavoidance, nutrient acquisition, host damage, biofilm formation and endotoxicity. The most defining characteristics of *Proteus* bacteria is the swarming phenomenon, a multicellular differentiation process of short rods to elongated swarmer cells. It allows the

population of bacteria to migrate in a coordinate manner on a solid surface. The role of short rods and elongated swarmer cells in the pathogenicity of *Proteus* bacteria is still not established. The elongated morphotype seems to be in the minority during infection. *Proteus* bacteria form a crystalline biofilm, particularly on the surface of catheters present in the urinary tract of patients. It contains struvite and hydroxyapatite crystals formed due to the action of urease. The results of the study on *Proteus* virulence and fitness factors will lead to a better understanding of infectious processes and will subsequently allow developing new methods of prevention and clinical treatment.

#### 5. References

- ALAMURI, P., MOBLEY, H.L.T. 2008. A novel autotransporter of uropathogenic *Proteus mirabilis* is both a cytotoxin and an agglutinin. *Mol. Microbiol.* 68: 997–1017.
- ALLISON, C., COLEMAN, N., JONES, P.L., HUGES, C. 1992. Ability of *Proteus mirabilis* to invade urothelial cells is coupled to motility and swarming differentiation. *Infect. Immun.* 60: 4740–4746.
- ALLISON, C., LAI, H.C., HUGHES, C. 1993. Cell differentiation of *Proteus mirabilis* is initiated by glutamine, a specific chemoattractant for swarming cells. *Mol. Microbiol.* 8: 53–60.
- ALTMAN, E., HARRISON, B.A., LATTA, R.K., LEE, K.K., KELLY, J.F., TIMBAULT, P. 2001. Galectin-3-mediated adherence of *Proteus mirabilis* to Madin-Darby canine kidney cells. *Biochem. Cell. Biol.* 79: 783–788.
- AQUILINI, E., AZEVEDO, J., JIMENEZ, N., BOUAMAMA, L., TOMAS, J.M., REGUE, M. 2010. Functional identification of the *Proteus mirabilis* core lipopolysaccharide biosynthetic genes. *J. Bacteriol.* 192: 4413–4424.
- BABICKA, D. 2001. Investigations of serological specificity of O-antigens of *Proteus vulgaris* O1, O2, O4 and O9 and selected biological features of this species. Ph.D. thesis. University of Lodz. (In Polish).
- BAHRANI, F.K., JOHNSON, D.E., ROBBINS, D. M., MOBLEY, H.L.T. 1991. *Proteus mirabilis* flagella and MR/P fimbriae: isolation, purification, N-terminal analysis, and serum antibody response following urinary tract infection. *Infect. Immun.* 59: 3574–3480.
- BELAS, R. 1994. Expression of multiple flagellin-encoding genes of *Proteus mirabilis*. *J. Bacteriol.* 176: 7169–7181.
- BELAS, R., MANOS, J., SUVANASUTHI, R. 2004. *Proteus mirabilis* ZapA metalloprotease degrades a broad spectrum of substrates, including antimicrobial peptides. *Infect. Immun.* 72: 5159–5167.
- BELAS, R., SUVANASUTHI, R. 2005. The ability of *Proteus mirabilis* to sense surfaces and regulate virulence gene expression involves FliL, a flagellar basal body protein. *J. Bacteriol.* 187: 6789–6803.
- BERLICKI, L., BOCHNO, M., GRABOWIECKA, A., BIALAS, A., KOSIKOWSKA, P., KAFARSKI, P. 2011. N-substituted aminomethanephosphonic and aminomethane-*P*-methylphosphinic acids as inhibitors of ureases. *Amino Acids*. DOI: 10.1007/s00726-011-0920-4.

- BEVERIDGE, T.J. 1999. Structures of Gram-negative cell walls and their derived membrane vesicles. *J. Bacteriol.* 181: 4725–4733.
- BIJLSMA, I.G., VAN DIJK, L., KUSTERS, J.G., GAASTRA, W. 1995. Nucleotide sequence of two fimbrial subunit genes, *pmpA* and *ucaA* from canine -uropathogenic *Proteus mirabilis* strains. *Microbiology.* 141: 1349–1357.
- BLOCH, J., LEMAIRE, X., LEGOUT, L., FERRIBY, D., YAZDANPANAH, Y., SENNEVILLE, E. 2010. Brain abscesses during *Proteus vulgaris* bacteremia. *Neurol. Sci.* 32: 661–663.
- CARSON, L., CATHCART, G.R., SCOTT, C.J., HOLLENBERG, M.D., WALKER, B., CERI, H., GILMORE B.F. 2011. Comprehensive inhibitory profiling of the *Proteus mirabilis* metalloprotease virulence factor ZapA (mirabilysin). *Biochemie.* 93:1824–1827.
- CARSON, L., GORMAN, S.P., GILMORE, B.F. 2010. The use of lytic bacteriophages in the prevention and eradication of biofilms of *Proteus mirabilis* and *Escherichia coli*. *FEMS Immunol. Med. Microbiol.* 59: 447–455.
- CHIPPENDALE, G.R., WARREN, J.W., TRIFILLIS, A.L., MOBLEY, H.L.T. 1994. Internalization of *Proteus mirabilis* by human renal epithelial cells. *Infect. Immun.* 62: 3115–3121.
- CHLABICZ, S., LESZCZYŃSKA, K., LUKAS, W., GUALCO, L., SCHITO, G., NABER, K.G. 2011. Uncomplicated lower urinary tract infections in females-clinical aspects, aetiology and antimicrobial resistance epidemiology. Results of the ARESC (Antimicrobial Resistance Epidemiological Survey on Cystitis) study in Poland on their implications for empiric therapy. *Przegl. Epidemiol.* 65: 345–351. (In Polish).
- COKER, C., POORE A., LI, X., MOBLEY, H.L.T. 2000. Pathogenesis of *Proteus mirabilis* urinary tract infections. *Microb. Infect.* 2: 1497–1505.
- DANIELS, R., VANDERLEYDEN, J., MICHIELS, J. 2004. Quorum sensing and swarming migration in bacteria. *FEMS Microbiol. Rev.* 28: 261–289.
- DONLAN, R.M., COSTERTON, J.W. 2002. Biofilms: survival mechanism of clinically relevant microorganisms. *Clin. Microbiol. Rev.* 15: 167–193.
- DOUFOR, A., FURNESS, R.B., HUGHES, C. 1998. Novel genes that upregulate the *Proteus mirabilis* *flhDC* master operon controlling flagellar biogenesis and swarming. *Mol. Microbiol.* 29: 741–751.
- DRECHSEL, H.A., THIECKEN, A., REISSBRODT, R., JUNG, G., WINKELMANN, G. 1993.  $\alpha$ -keto acids are novel siderophores in the genera *Proteus*, *Providencia* and *Morganella* and are produced by amino acid deaminases. *J. Bacteriol.* 175: 2727–2733.
- DRZEWIECKA, D., ARBATSKY, N.P., SHASHKOV, A.S., STĄCZEK, P., KNIREL, Y.A., SIDORCZYK, Z. 2008. Structure and serological properties of the O-antigen of two clinical *Proteus mirabilis* strains classified into a new *Proteus* O77 serogroup. *FEMS Immunol. Med. Microbiol.* 54: 185–194.
- DRZEWIECKA, D., ARBATSKY, N.P., STĄCZEK, P., SHASHKOV, A.S., KNIREL, Y.A., SIDORCZYK, Z. 2010. Structure and serological studies of the O-polysaccharide of strains from a newly created *Proteus* O78 serogroup prevalent in Polish patients. *FEMS Immunol. Med. Microbiol.* 58: 269–276.
- DRZEWIECKA, D., SIDORCZYK, Z. 2005. Characterization of *Proteus penneri* species – human opportunistic pathogens. *Post. Mikrobiol.* 44: 113–126. (In Polish).
- DRZEWIECKA, D., ZYCH, K., SIDORCZYK, Z. 2004. Characterization and serological classification of a collection of *Proteus penneri* clinical strains. *Arch. Immunol. Therap. Exp.* 52:121–128.
- DUMANSKI, A.J., HEDELIN, H., EDIN-LILJEGREN, A., BEACHEMIN, D., MCLEAN, R.J.C. 1994. Unique ability of *Proteus mirabilis* capsule to enhance mineral growth in infectious urinary calculi. *Infect. Immun.* 62: 2998–3003.
- FLANNERY, E., MODY, L., MOBLEY, H.L.T. 2009. Identification of a modular pathogenicity island that is widespread among urease-producing uropathogens and shares features with diverse group of mobile elements. *Infect. Immun.* 77: 4887–4894.
- FUJIHARA, M., OBARA, H., WATANABE, Y., ONO, H.K., SASAKI J., GORYO, M., HARASAWA, R. 2011. Acidic environments induce differentiation of *Proteus mirabilis* into swarmer morphotypes. *Microbiol. Immunol.* 55: 489–493.
- GIAMMANCO, G.M., GRIMONT, P.A., GRIMONT, F., LEFÈVRE, M., GIAMMANCO, G., PIGNATO, S. 2011. Phylogenetic analysis of the genera *Proteus*, *Morganella*, *Providencia* by comparison of *rpoB* gene sequence of type and clinical strains suggests reclassification of *Proteus myxofaciens* in a new genus, *Cosenzaea* gen. nov., as *Cosenzaea myxofaciens* comb. nov. *Int. J. Syst. Evol. Microbiol.* 61:1638–1644.
- GYGI, D., RAHMAN, M.M., LAI, H.C., CARLSON, R., GUARD-PETTER, J., HUGHES, C. 1995. A cell surface polysaccharide that facilitates rapid population migration by differentiated swarm cells of *Proteus mirabilis*. *Mol. Microbiol.* 17: 1167–1175.
- HART, B.R., BLUMENTHAL, R.M. 2010. Unexpected coregulator range in the global regulator Lrp of *Escherichia coli* and *Proteus mirabilis*. *J. Bacteriol.* 193: 1054–1064.

- HART, B.R., MISHRA, P.K., LINTER, R.E., HINERMAN, J.M., HERR, A.B., BLUMENTHAL, R.M. 2011. Recognition of DNA by the helix-turn-helix global regulatory protein Lrp is modulated by the amino terminus. *J. Bacteriol.* 193:3794–3803.
- HATT, J.K., RATHER, P.N. 2008. Characterization of novel gene, *wosA*, regulating FlhDC expression in *Proteus mirabilis*. *J. Bacteriol.* 190: 1946–1955.
- HAY, N.A., TIPPER, D., GYGI, D., HUGHES, C. 1997. A nonswarming mutant of *Proteus mirabilis* lacks the Lrp global transcriptional regulator. *J. Bacteriol.* 179: 4741–4746.
- HIMPSI, S.D., PEARSON, M.M., AREWANG, C.J., NUSCA T.D., SHERMAN D.H., MOBLEY, H.L.T. 2010. Proteobactin and a yersiniabactin-related siderophore mediate iron acquisition in *Proteus mirabilis*. *Molec. Microbiol.* 78: 138–157.
- HOLST, O. 1999. Chemical structure of the core region of lipopolysaccharides. In: H. BRADE, S.M. OPAL, S.N. VOGEL, D.C. MORRISON (eds). *Endotoxin in health and disease*. Marcel Dekker, Inc. New York, Basel, pp. 115–154.
- JACOBSEN, S.M., STICKLER D.J., MOBLEY, H.L.T., SHIRTLIFF, M.E. 2008. Complicated catheter-associated urinary tract infections due to *Escherichia coli* and *Proteus mirabilis*. *Clin. Microbiol. Rev.* 21: 26–59.
- JANDA, J.M., ABBOT, S.L. 2006. *The Enterobacteriaceae*. ASM Press. Washington. 233–259.
- JANSEN, A.M., LOCKATELL, C.V., JOHNSON, D.E., MOBLEY, H.L.T. 2004. Mannose-resistant *Proteus*-like fimbriae are produced by most *Proteus mirabilis* strains infecting the urinary tract, dictate the *in vivo* localization of bacteria, and contribute to biofilm formation. *Infect. Immun.* 72: 7292–7305.
- JANSEN, B.V., LOCKATELL, C.V., JOHNSON, D.E., MOBLEY, H.L.T. 2003. Visualisation of *Proteus mirabilis* morphotypes in the urinary tract: the elongated swarmer cell is rarely observed in ascending urinary tract infection. *Infect. Immun.* 71: 3607–3613.
- JONES, B.D., LOCKATELL, C.V., JOHNSON, D.E., WARREN, J.W., MOBLEY, H.L.T. 1990. Construction of urease-negative mutant of *Proteus mirabilis*: analysis of virulence in mouse model of ascending urinary tract infection. *Infect. Immun.* 58: 1120–1123.
- JONES, S.M., YERLY, J., HU, Y., CERI, H., MARTINUZZI, R. 2007. Structure of *Proteus mirabilis* biofilm grown in artificial urine and standard laboratory media. *FEMS Microbiol. Lett.* 268:16021.
- JONES, S.M., DANG, T., MARTINUZZI, R. 2009. Use of quorum sensing antagonists to deter the formation of crystalline *Proteus mirabilis* biofilm. *Int. J. Antimicrob. Agen.* 34: 360–364.
- KACA, W., ARABSKI, M., FUDAŁA, R., HOLMSTROM, E., SJOHOLM, A., WEINTRAUB, A., FUTOMA-KOLOCH., B., BUGLA-PŁOSKOŃSKA, G., DOROSZKIEWICZ, W. 2009. Human complement activation by smooth and rough *Proteus mirabilis* lipopolysaccharides. *Arch. Immunol. Ther. Exp.* 57: 383–391.
- KACA, W., GLENSKA, J., LACHOWICZ, L., GRABOWSKI, S., BRAUNER, A., KWINKOWSKI, M. 2011. Serotyping of *Proteus mirabilis* clinical strains based on lipopolysaccharide, O-polysaccharide and core oligosaccharide structures. *Biochemistry (Moscow)*. 76: 851–861.
- KACA, W., UJAZDA, E. 1998. Complement activation by bacterial endotoxin. *Post. Mikrobiol.* 37: 421–427. (In Polish).
- KALRA, A., COOLEY, C., TSIGRELIS, C. 2011. Treatment of endocarditis due to *Proteus* species: a literature review. *Int. J. Infect. Dis.* 15: 222–225.
- KNIREL, Y.A., PEREPELOV, A.V., KONDAKOVA, A., SENCHEKOVA, S.N., SIDORCZYK, Z., RÓZALSKI, A., KACA, W. 2011. Structure and serology of O-antigens as the basis for classification of *Proteus* strains. *Inn. Immun.* 17: 70–96.
- KORONAKIS, V., CROSS, M., SENIOR, B., KORONAKIS, E., HUGHES, C. 1987. The secreted hemolysins of *Proteus mirabilis*, *Proteus vulgaris*, and *Morganella morganii* are genetically related to each other and to the alpha-hemolysin of *Escherichia coli*. *J. Bacteriol.* 169: 1509–1515.
- KWIECIŃSKA-PIRÓG, J. 2010. Estimation of production of biofilm by *Proteus mirabilis* bacilli. Ph.D. thesis. Nicolaus Copernicus University of Toruń. Collegium Medicum in Bydgoszcz. (In Polish).
- KWIL, I., BABICKA, D., STĄCZEK, P., RÓZALSKI, A. 2002. Applying the PCR metod for searching genes encoding main structural fimbrial proteins. In: 5<sup>th</sup> Conference on Molecular biology in diagnostics of infectious disease and biotechnology. Warsaw, SGGW Publisher. 88–91.
- KWIL, I. 2003. Investigations of selected virulence factors of *Proteus penneri* strains. Ph.D. thesis. University of Lodz. (In Polish).
- KWIL, I., BIEGAŃSKA, H., OBAREK, K., RÓZALSKI, A. 2006. Expression of selected virulence factors among new bacterial species from the genus *Proteus*: *P. hauseri* and *Proteus* genomospecies 4, 5, 6. In: 9<sup>th</sup> Conference on molecular biology in diagnostics of infectious disease and biotechnology. Warsaw, SGGW Publisher. 93–96.
- LANE, M.C., PEARSON, M.M., SIMMS, A.N., MOBLEY, H.L.T. 2009. Oxygen-limiting conditions enrich for fimbriate cells of uropathogenic *Proteus mirabilis* and *Escherichia coli*. *J. Bacteriol.* 191: 1382–1392.

- LEE, K.K., HARRISON, B.A., LATTA, R., ALTMAN, E. 2000. The binding of *Proteus mirabilis* nonagglutinating fimbriae to ganglio-series asialoglycolipids and lactosyl ceramide. *Can. J. Microbiol.* 46: 961–966.
- LI, X., JOHNSON, D.E., MOBLEY, H.L.T. 1999. Requirement of MrpH for mannose-resistant *Proteus*-like fimbria-mediated hemagglutination by *Proteus mirabilis*. *Infect. Immun.* 67: 2822–2833.
- LI, X., LOCKATELL, C.V., JOHNSON, D.E., MOBLEY, H.L.T. 2002a. Identification of MrpI as the sole recombinase that regulates the phase variation of MR/P fimbria, a bladder colonization factor of uropathogenic *Proteus mirabilis*. *Mol. Microbiol.* 45: 865–874.
- LI, X., ZHAO, H., LOCKATELL, C.V., DRACHENBERG, C.B., JOHNSON, D.E., MOBLEY, H.L.T. 2002b. Visualization of *Proteus mirabilis* within the matrix of urease-induced bladder stones during experimental urinary tract infection. *Infect. Immun.* 70: 389–394.
- LIAW, S.J., LAI, H.C., HO, S.W., LUH, K.T., WANG, W.B. 2003. Role of RsmA in the regulation of swarming motility and virulence factor expression in *Proteus mirabilis*. *J. Med. Microbiol.* 52: 19–28.
- LIAW, S.J., LAI, H.C., WANG, W.B. 2004. Modulation of swarming and virulence by fatty acids through the RsbA protein in *Proteus mirabilis*. *Infect. Immun.* 72: 6836–6845.
- LOOMES, L.M., SENIOR, B.W., KERR, M.A. 1990. A proteolytic enzyme secreted by *Proteus mirabilis* degrades immunoglobulins of the immunoglobulin A1 (IgA1), IgA2, and IgG isotypes. *Infect. Immun.* 58: 1970–1985.
- ŁUKASIEWICZ, J., ŁUGOWSKI, C. 2003. Biological activities of lipopolysaccharide. *Post. Hig. Med. Doświad.* 57: 33–53. (In Polish).
- MALIC, W., WATTERS, M.G., BASIL, L., STICKLER, D.J., WILLIAMS, D.M. 2011. Development of an “early warning” sensor for encrustation of urinary catheters following *Proteus* infection. *J. Biomed. Mater. Res. B Appl. Biomater.* DOI: 10.1002/jbm.b.31930.
- MASSAD, G., ZHAO, H., MOBLEY, H.L.T. 1995. *Proteus mirabilis* amino acid deaminase: cloning, nucleotide sequence, and characterization of *aad*. *J. Bacteriol.* 177: 5878–5883.
- MIELNIK, G., DOROSZKIEWICZ, W., KORZENIOWSKA-KOWAL, A. 2004. External structures of Gram-negative bacteria and bactericidal activity of complement. *Post. Mikrobiol.* 43: 39–57. (In Polish).
- MOBLEY, H.L.T. 1996. Virulence of *Proteus mirabilis*. In: H.L.T. MOBLEY, J.W. WARREN (eds.) *Urinary tract infections, molecular pathogenesis and clinical management*. ASM Press, Washington DC, pp. 245–269.
- MOBLEY, H.L.T., BELAS, R., LOCKATELL, V., CHIPPENADALE, G., TRIFILLIS A.L., JOHNSON, D.E., WARREN, J.W. 1996. Construction of a flagellum-negative mutant of *Proteus mirabilis*: effect on internalization by human renal epithelial cells and virulence in a mouse model of ascending urinary tract infection. *Infect. Immun.* 64: 5332–5340.
- MOBLEY, H.L.T., ISLAND, M.D., HAUSINGER, R.P. 1995. Molecular biology of microbial ureases. *Microbiol. Rev.* 59: 451–480.
- MORGENSTEIN, R.M., SZOSTEK, B., RATHER, P.N. 2010. Regulation of gene expression during swarmer cell differentiation in *Proteus mirabilis*. *FEMS Microbiol. Rev.* 34: 753–763.
- MORYL, M. 2010. Investigations of biofilm formation by *Proteus mirabilis* and its sensitivity to antibacterial agents. Ph.D. Thesis. University of Lodz. (In Polish).
- MORYL, M., TORZEWSKA, A., RÓZALSKI, A. 2008. Substances affecting *Proteus mirabilis* biofilm. In: Conference “Vaccines: Advances in plant and microbial biotechnology, infectious immunity and cancer therapy. Warsaw, SGW Publisher. 179–181.
- NIELUBOWICZ, G., MOBLEY, H.L.T. 2010. Host-pathogen interactions in urinary tract infections. *Nature Rev. Urology* 7: 430–441.
- O’ HARA, C., BRENNER, F.W., MILLER, J.M. 2000. Classification, identification, and clinical significance of *Proteus*, *Providencia*, and *Morganella*. *Clin. Microbiol. Rev.* 13: 534–546.
- OKIMOTO, N., HAYASHI, T., ISHIGA, M., NANBA, F., KISHIMOTO, M., YAGI, S., KURIHARA, T., ASOKA, N., TAMADA, S. 2010. Clinical features of *Proteus mirabilis* pneumonia. *J. Infect. Chemother.* 16: 364–366.
- PALUSIAK, A., SIDORCZYK, Z. 2009. Serological characterization of the core region of lipopolysaccharides of rough *Proteus* sp. strains. *Arch. Immunol. Ther. Exp.* 57: 303–310.
- PALUSIAK, A., SIDORCZYK, Z. 2010. Characterization of epitope specificity of *Proteus penneri* 7. *ABP.* 57: 529–532.
- PEARSON, M.M., MOBLEY, H.L.T. 2008. Repression motility during fimbrial expression; identification of 14 *mrpJ* gene paralogues in *Proteus mirabilis*. *Mol. Microbiol.* 69: 548–558.
- PERSON, M.M. *et al.* 2008. Complete genome sequence of uropathogenic *Proteus mirabilis*, a master of both adherence and motility. *J. Bacteriol.* 190: 4027–4037.

- PHAN, V., BELAS, R., GILMORE, B.F., CERI, H. 2008. ZapA, a virulence factor in a rat model of *Proteus mirabilis*-induced acute and chronic prostatitis. *Infect. Immun.* 76: 4859–4864.
- PIECHOTA, J., PRYWER, J., TORZEWSKA, A. 2012. *Ab initio* prediction of structural and elastic properties of struvite: contribution to urinary stone research. *Comput. Meth. Biochem. Engin.* 15: 1329–1336.
- PRYWER, J., TORZEWSKA, A. 2009. Bacterially induced struvite growth from synthetic urine: experimental and theoretical characterization of crystal morphology. *Cryst. Growth Design.* 9: 3538–3543.
- PRYWER, J., TORZEWSKA, A. 2010. Biomineralization of struvite crystals by *Proteus mirabilis* from artificial urine and their mesoscopic structure. *Cryst. Res. Technol.* 45: 1283–1289.
- PRYWER, J., TORZEWSKA, A. 2012. Effect of curcumin against *Proteus mirabilis* during crystallization of struvite from artificial urine. *Evidence Base Complementary and Alternative Medicin.* 862794: 1–7.
- RADZIEJEWSKA-LEBRECHT, J., MAYER, H. 1989. The core region of *Proteus mirabilis* R110/1959 lipopolysaccharide. *Eur. J. Biochem.* 183: 573–581.
- RAETZ, C.R., WHITFIELD, C. 2002. LIPOPOLYSACCHARIDE ENDOTOXIN. *ANN. REV. BIOCHEM.* 71: 635–700.
- RATHER, P.N. 2005. Swarmer cell differentiation in *Proteus mirabilis*. *Environm. Microbiol.* 7: 1065–1073.
- ROCHA, S.P.D., ELIAS, W.P., CIANCARULLO, A.M., MENEZES M.A., NARA, J.M., PIAZZA, R.F.M., SILVA, M.R.L., MOREIRA, C.G., PELAYO, J.S. 2007a. Aggregative adherence of uropathogenic *Proteus mirabilis* to cultured epithelial cells. *FEMS Immunol. Med. Microbiol.* 51: 319–326.
- ROCHA, S.P.D., ELIAS, W.P., PELAYO, J.S. 2007b. Fimbriae of uropathogenic *Proteus mirabilis*. *FEMS Immunol. Med. Microbiol.* 51: 1–7.
- ROMANOWSKI, Z., KEMPISTY, P., PRYWER, J., KRUKOWSKI, S., TORZEWSKA A. 2010. Density functional theory determination of structural and electronic properties of struvite: *J. Phys. Chem.* 114: 7800–7808.
- RÓZALSKI, A. 2002. Molecular basis of the pathogenicity of *Proteus* bacteria. *Adv. Clin. Exp. Med.* 11: 3–18.
- RÓZALSKI, A. 2004. Lipopolysaccharide and others virulence factors of *Proteus* bacteria. *Post. Mikrobiol.* 43: 409–431. (In Polish).
- RÓZALSKI, A. 2008. Lipopolysaccharide (LPS, endotoxin) of *Proteus* bacteria – chemical structure, serological specificity and the role in pathogenicity. *Folia Biolog. Oecolog.* 4: 5–24.
- RÓZALSKI, A., DŁUGOŃSKA, H., KOTELKO, K. 1986. Cell invasiveness of *Proteus mirabilis* and *Proteus vulgaris* strains. *Arch. Immunol. Ther. Exp.* 34: 505–511.
- RÓZALSKI, A., KOTELKO, K. 1987. Hemolytic activity and invasiveness in strains of *Proteus penneri*. *J. Clin. Microbiol.* 25: 1094–1096.
- RÓZALSKI, A., SIDORCZYK, Z., KOTELKO, K. 1997. Potential virulence factors of *Proteus* bacilli. *Microbiol. Mol. Biol. Rev.* 61: 65–89.
- RÓZALSKI, A., STĄCZEK, P. 2010. *Proteus*. In: D. LIU, (ed.) *Molecular detection of foodborne pathogens*. CRS Press, Taylor and Francis Group. Boca Raton, p. 417–430.
- RÓZALSKI, A., STĄCZEK, P. 2011. *Proteus*. In: *Molecular detection of human bacterial pathogens*. In: D. Liu (ed.), CRC Press, Taylor and Francis Group. Boca Raton, p. 981–996.
- RÓZALSKI, A., TORZEWSKA, A., BARTODZIEJSKA B., BABICKA D., KWIL, I., PEREPELOV, A.V., KONDAKOVA, A.N., SENCHENKOVA, S.N., KNIREL, Y.A., VINOGRADOV, E.V. 2002. Chemical structure, antigenic specificity, the role in the pathogenicity of lipopolysaccharide (LPS, endotoxin) one the *Proteus vulgaris* bacteria's example. *Wiadomości Chemiczne.* 56: 585–604. (In Polish).
- SCAVONE, P., UMPIERREZ, A., MASKELL, D.J., ZUNINO, P. 2011. Nasal immunization with attenuated *Salmonella* Thyphimurium expressing and MrpA-TetC fusion protein significantly reduces *Proteus mirabilis* colonization in mouse urinary tract. *J. Med. Microbiol.* 60: 899–904.
- SCHLAPP, G., SCAVONE, P., ZUNINO, P., HARTEL, S. 2011. Development of 3D architecture of uropathogenic *Proteus mirabilis* bath culture biofilms – A quantitative confocal microscopy approach. *J. Microbiol. Meth.* 87: 234–240.
- SIDORCZYK, Z., ZAHNINGER, U., RIETSCHER, E.T. 1983. Chemical structure of the lipid A component of the lipopolysaccharide of *Proteus mirabilis* Re mutant. *Eur. J. Biochem.* 137: 15–22.
- SOSA, V., ZUNINO, P. 2009. Effect of *Ibicella lutea* on uropathogenic *Proteus mirabilis* growth, virulence and biofilm formation. *J. Infect. Dev. Ctries.* 3: 762–770.
- STAŃKOWSKA, D., CZERWONKA, G., RÓZALSKA, S., GROSICKA, M., DZIADEK, J., KACA, W. 2012. Influence of quorum sensing signal molecules on biofilm formation in *Proteus mirabilis* O18. *Folia Microbiol.* 57: 53–60.
- STAŃKOWSKA, D., KWINKOWSKI, M., KACA, W. 2008. Quantification of *Proteus mirabilis* virulence factors and modulation by acylated homoserine lactones. *J. Microbiol. Immunol. Infect.* 41: 243–253.



- STICKLER, D.J., 2008. Bacterial biofilms in patients with indwelling urinary catheters. *Nature Clin. Pract. Urology*. 5: 598–607.
- STICKLER, D.J., FENELEY, R.C.L. 2010. The encrustation and blockage of long-term indwelling bladder catheters: a way forward in prevention and control. *Spinal Cord*. 48: 784–790.
- SWIHART, K.G., WELCH, R.A. 1990. Cytotoxic activity of *Proteus* hemolysin HpmA. *Infect. Immun.* 58: 1861–1869.
- TORZEWSKA, A., RÓZALSKI, A. 2009. Studies on infection-induced urinary calculi formation in the presence of glycosaminoglycans. X Jubilee Conference "Molecular biology in diagnostic of infectious diseases and biotechnology, Warsaw University of Life Sciences - SGGW, 2009. Scientific materials, 179–182.
- TORZEWSKA, A., STĄCZEK, P., RÓZALSKI, A. 2003. The crystallization of urine mineral components may depend on the chemical nature of *Proteus* endotoxin polysaccharides. *J. Medical. Microbiol.* 52: 471–477.
- TORZEWSKA, A., ZIELIŃSKA, E., ŚWIDER, S., RÓZALSKI, A. 2010. Influence of curcumin on *Proteus mirabilis* pathogenicity. *Sepsis*. 4: 276–277.
- UPHOFF, T.S., WELCH, R.A. 1990. Nucleotide sequencing of the *Proteus mirabilis* calcium independent hemolysin genes (*hpmA* and *hpmB*) reveals sequence similarity with *Serratia marcescens* hemolysin genes (*shlA* and *shlB*). *J. Bacteriol.* 172: 1206–1216.
- VERSTRAETEN, N., BREKANEN, K., DEBKUMARI, B., FAUVART, M., FRANSAER, J., VERMANT J., MICHELS, J. 2008. Living on a surface: swarming and biofilm formation. *Trends Microbiol.* 16: 496–506.
- VINOGRADOV, E. 2011. Structure of the core part of the lipopolysaccharide from *Proteus mirabilis* genomic strain HI4320. *Biochemistry (Moscow)*. 76: 803–807.
- VINOGRADOV, E.V., SIDORCZYK, Z., KNIREL, Y.A. 2002. Structure of lipopolysaccharide core region of the genus *Proteus*. *Aust. J. Chem.* 55: 61–67.
- VINOGRADOV, E.V., THOMAS-OATES, E., BRADE, H., HOLST, O. 1994. Structural investigations of the lipopolysaccharide from *Proteus mirabilis* R45 (Re-chemotype). *J. End. Res.* 1: 199–206.
- WANG, Q., TORZEWSKA, A., RUAN, X., WANG, X., RÓZALSKI, A., SHAO, Z., GUO, X., FENG, L., WANG, L. 2010. Molecular and genetic analyses of the putative *Proteus* O-antigen gene locus. *App. Env. Microbiol.* 76: 5471–5480.
- WARREN, J.W. 1996. Clinical presentations and epidemiology of urinary tract infections. In: H.L.T. MOBLEY, J.W. WARREN (eds.) *Urinary tract infections, molecular pathogenesis and clinical management*. ASM Press, Washington DC, p. 2–28.
- WRAY, S.K., HULL, S.I., COOK, R.G., BARRISH, J., HULL, R.A. 1986. Identification and characterization of a uroepithelial cell adhesion from a uropathogenic isolate of *Proteus mirabilis*. *Infect. Immun.* 54: 43–49.
- ZUNINO, P., GEYMONT, L., ALLEN, A.G., LEGNANI-FAJARDO, C., MASKELL, D.J. 2000. Virulence of *Proteus mirabilis atf* isogenic mutant is not impaired in a mouse model ascending urinary tract infection. *FEMS Immunol. Med. Microbiol.* 29: 137–143.
- ZUNINO, P., PICCINI, C., LEGNANI-FAJARDO, C. 1994. Flagellate and non flagellate *Proteus mirabilis* in the development of experimental urinary tract infection. *Microb. Path.* 16: 379–385.
- ZUNINO, P., SOSA, V., SHLAPP, G., ALLEN, A.G., PRESTON A., MASKELL, D.J. 2007. Mannose-resistant *Proteus*-like and *Proteus mirabilis* fimbriae have specific and additive roles in *P. mirabilis* urinary tract infections. *FEMS Immunol. Med. Microbiol.* 51: 125–133.